

temperature range but it eventually kill micro-organism that are required for optimum quantity of gas production. In addition, thermophilic conditions suitable in cold climate region or winter. Hence in our region where purpose of study to use solar based biogas plant. Mesophilic (35 °C-45 °C) temperature ranges were suitable to maintain temperature.

REFERENCES

- Fazenda, M.L., et al. 2008. Submerged culture fermentation of "higher fungi": the macrofungal. *Advances in applied microbiology*, 63, 33-103.
- Harikishan, S., Sung, S. 2003. Cattle waste treatment and Class A biosolid production using temperature-phased anaerobic digester. *Advances in Environmental Research*, 7 (3), 20, 701-706.
- Li, R., et al. 2009. Anaerobic co-digestion of kitchen waste and cattle manure for methane production. *Energy Sources, Part A: Recovery, Utilization, and Environmental Effects*, 20, 1848-1856.
- Mata-Alvarez, J., Mace, S., Llabres, P., 2000. Anaerobic digestion of organic solid wastes. An overview of research achievements and perspectives. *Bioresource technology*, 1, 3-16.
- Najafpour, G. 2007. *Bioreactor Design*, 15, 142-169.
- Nawaz, A., Hussain, S., Waqas, M.S., Rasheed, H., Ali, S., Iqbal, M.M., Yasmeen, Z., Waqas, M.M., 2020. Health risk assessment of heavy metals due to untreated wastewater irrigated vegetables. *Big data in water resources engineering (BDWRE)*, 1 (1), 01-03.
- Panwar, N., et al. 2011. Role of renewable energy sources in environmental protection: A review. *Renewable and Sustainable Energy Reviews*, 15,1513-1524.
- Singh, J., Kaushik, N., Biswas, S. 2014. Bioreactors–technology & design analysis. *The SciTech Journal*, 1, 28-36.
- Weiland, P., 2010. Biogas production: current state and perspectives. *Applied microbiology and biotechnology*, 4, 849-860.

